

The Role of Boundary Conditions in the Real-Space Renormalization Group ^{*}

Miguel A. Martín-Delgado[†] and Germán Sierra[‡]

[†]*Departamento de Física Teórica I
Universidad Complutense. 28040-Madrid, Spain*
[‡]*Instituto de Matemáticas y Física Fundamental. C.S.I.C.
Serrano 123, 28006-Madrid, Spain*

February 1, 2008

Abstract

We show that the failure of the real-space RG method in the 1D tight-binding model is not intrinsic to the method as considered so far but depends on the choice of boundary conditions. For fixed BC's the failure does happen. For free BC's we present a new analytical block RG-method which gives the exact ground state of the model and the correct $1/N^2$ -law for the energy of the first excited state in the large N (size)-limit. We also give a reconstruction method for the wave-functions of the excited states.

PACS numbers: 05.50.+q, 71.10.+x, 64.60.Ak

^{*}Work partly supported by CICYT under contracts AEN93-0776 (M.A.M.-D.) and PB92-109 , European Community Grant ERBCHRXCT920069 (G.S.).

1 Introduction

Real space Renormalization Group (RG) methods originated from the study of the Kondo problem by Wilson [1]. It was clear from the beginning that one could not hope to achieve the accuracy Wilson obtained for the Kondo problem when dealing with more complicated many-body quantum Hamiltonians. The key difference is that in the Kondo model there exists a *recursion relation* for Hamiltonians at each step of the RG-elimination of degrees of freedom. Schematically,

$$H_{N+1} = H_N + \text{hopping boundary term} \quad (1)$$

$$H_{N+1} = R(H_N) \quad (2)$$

where R is the RG transformation.

The existence of such recursion relation facilitates enormously the work, but as it happens it is specific of *impurity problems*.

From the numerical point of view, the Block Renormalization Group (BRG) procedure proved to be not fully reliable in the past particularly in comparison with other numerical approaches, such as the Quantum MonteCarlo method which were being developed at the same time. This was one of the reasons why the BRG methods remained undeveloped during the '80's until the beginning of the '90's when they are making a comeback as one of the most powerful numerical tools when dealing with zero temperature properties of many-body systems, a situation where the Quantum MonteCarlo methods happen to be particularly badly behaved as far as fermionic systems is concerned [2].

As it happens, the BRG gives a good qualitative picture of many properties exhibited by quantum lattice Hamiltonians: Fixed points, RG-flow, phases of the system etc. as well as good quantitative results for some properties such as ground state energy and others [3], [4], [5], [6]. For a review on the Block RG method see [7] and chapter 11 of reference [8]. However in some important instances the BRG method is off the correct values of critical exponents by a sensible amount.

The first advance in trying to understand the sometimes bad numerical performance of the BRG methods came in the understanding of the effect of *boundary conditions* (BC) on the standard RG procedure [9].

White and Noack [9] pointed out that the standard BRG approach of neglecting all connections to the neighbouring blocks during the diagonalization of the block Hamiltonian H_B introduces large errors which cannot be corrected by any reasonable increase in the number of states kept. Moreover, in order to isolate the origin of this problem they study an extremely simple model: a free particle in a 1D lattice. As a matter of fact, it was Wilson [10] who pointed out the importance of understanding real-space RG in the context of this simple tight-binding model where the standard BRG clearly fails as we are going to show.

The reason for this failure can be traced back to the importance of the boundary conditions in diagonalizing the states of a given block Hamiltonian H_B in which the lattice is decomposed into. Notice that in this fashion we are isolating a given block from the rest of the lattice and this applies a *particular* BC to the block. However, the block is not truly isolated! A statement which is the more relevant the more strongly correlated is the system under consideration. Thus, if the rest of the lattice were there it would apply different BC's to the boundaries of the block. This in turn makes the standard block-diagonalization conceptually not faithfully suited to account for the interaction with the rest of the lattice.

Once the origin of the problem is brought about the solution is also apparent: devise a method to change the boundary conditions in the block in order to mimic the interaction with the rest of the lattice. This is called the Combination of Boundary Conditions (CBC) method which yields very good numerical results. This method has not yet been generalized to interacting systems. However in reference [11] an alternative approach is proposed under the name of Density Matrix Renormalization

Group (DMRG) which applies to more general situations and also produces quite accurate results. Despite of the impressive numerical accuracy of these methods there is not a clear understanding of why they perform so well or whether they would be valid in dimensions higher than one.

2 Boundary Conditions and Real-Space RG

In this letter we want to reconsider again the role of BC's in the real space RG method for the case of a single-particle problem in a box. The continuum version of this Hamiltonian is simply $H = -\frac{\partial^2}{\partial x^2}$. We shall consider open chains with two types of BC's at the ends:

$$\text{Fixed BC's: } \psi(0) = \psi(L) = 0 \quad (3)$$

$$\text{Free BC's: } \frac{\partial\psi}{\partial x}(0) = \frac{\partial\psi}{\partial x}(L) = 0 \quad (4)$$

The lattice version of H for each type of BC's is given as follows:

$$H_{Fixed} = \begin{pmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & & \\ & & & \ddots & \\ & & & & 2 & -1 \\ & & & & -1 & 2 \end{pmatrix} \quad (5)$$

$$H_{Free} = \begin{pmatrix} 1 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & & \\ & & & \ddots & \\ & & & & 2 & -1 \\ & & & & -1 & 1 \end{pmatrix} \quad (6)$$

The only difference between H_{Fixed} and H_{Free} appear at the first and last diagonal entry ($2 \leftrightarrow 1$). The exact solution of (5) and (6) is very well-known and we give it for completeness:

$$\text{Fixed BC's: } \psi_n(j) = N_n^{Fx} \sin \frac{\pi(n+1)}{N+1} j, \quad E_n = 4 \sin^2\left(\frac{\pi(n+1)}{2(N+1)}\right) \quad (7)$$

$$\text{Free BC's: } \psi_n(j) = N_n^{Fr} \cos \frac{\pi n}{N} \left(j - \frac{1}{2}\right), \quad E_n = 4 \sin^2\left(\frac{\pi n}{2N}\right) \quad (8)$$

$$j = 1, 2, \dots, N; \quad n = 0, 1, \dots, N-1.$$

where the N'_n s are normalization constants and N is the number of sites of the chain.

Before getting into the problem of the renormalization of these Hamiltonians, it is worth to pointing out another physical realization of H_{Free} . Consider the following Hamiltonian for the anisotropic Heisenberg model in an open chain of N sites,

$$H_{\Delta} = -\frac{1}{2} \sum_{j=1}^{N-1} (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \Delta(\sigma_j^z \sigma_{j+1}^z - 1)) \quad (9)$$

where $\Delta = +1$ ($\Delta = -1$) corresponds to the Ferromagnetic case (Antiferromagnetic case). When $\Delta > 1$, the ground state of H_{Δ} is the Ferromagnetic state $|F\rangle$ given by:

$$|F\rangle = |\uparrow\uparrow\ldots\uparrow\rangle \quad (10)$$

$$H_\Delta|F\rangle = 0 \quad (11)$$

Now, let us denote by $|x\rangle$ the state which is obtained from the Ferromagnetic state $|F\rangle$ when one spin is overturned at the position x of the chain. The subspace spanned by the $|x\rangle$ states is the 1 magnon subspace of the Ferromagnetic model. It is readily checked that the restriction of the H_Δ Ferromagnetic Heisenberg Hamiltonian takes the following form:

$$H_{Magnon}(\Delta) = \begin{pmatrix} 1 & -1 & & & & \\ -1 & 2\Delta & -1 & & & \\ & -1 & 2\Delta & & & \\ & & & \ddots & & \\ & & & & 2\Delta & -1 \\ & & & & -1 & 1 \end{pmatrix} \quad (12)$$

If $\Delta = +1$ it precisely coincides with H_{Free} :

$$H_{Magnon}(\Delta = +1) = H_{Free} \quad (13)$$

Thus we arrive at the following mapping: *A simple magnon above a ferromagnetic background satisfies Free BC's.* We shall come back to this connection at the conclusions for it opens the door to the introduction of interactions in the model.

Now let us get to the problem of renormalizing the tight-binding Hamiltonians (5)-(6).

In figures 1 and 2 we show the ground state and first excited states of the chain with fixed and free BC's.

It is clear from Fig.1 that *a standard Block RG method is not appropriate to study the ground state of fixed BC's since this state is non-homogeneous while the block truncation does not take into account this fact.*

Each piece of the ground state within each block satisfies BC's which vary from block to block. This is the motivation of reference [9] to consider different BC's in the block method, yielding quite accurate results as can be seen from Table 1 for comparison of the CBC method and the standard method.

We observe that the standard RG method performs rather poorly as compared to the CBC method which yields quite the exact results.

The other alternative to the CBC method is the Density Matrix RG method which can be phrased by saying that the rest of the chain produces on every block the appropriate BC's to be applied to its ends, and it has the virtue that can be generalized to other models, something which is not the case as for the CBC method.

On the other hand, the ground state of H_{Free} is an homogeneous state (see Fig.2) which in turn suggests that a standard RG analysis may work for this type of BC's. We shall show that this is indeed the case if the RG procedure is properly defined. The key of our RG-prescription is to notice that H_{Free} has a geometrical meaning: *H_{Free} is the incidence matrix of the graph in Fig.3*, and it is called minus the discrete laplacian $-\Delta$ of that graph. Notice that H_{Fixed} has not such geometrical interpretation, in fact, it coincides with the Dynkin diagram of the algebra A_N .

Based on this observation the Kadanoff blocking is nothing but the breaking of the graph into N/n_s disconnected graphs of n_s sites each. We shall choose $n_s = 3$ in our later computation as shown in Fig. 4.

The previous geometrical interpretation of H_{Free} suggests that we choose the block Hamiltonian H_B to be the incidence matrix of the disconnected graph in Fig.4, namely,

$$H_B = \begin{pmatrix} 1 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 1 & & \\ & & & 1 & -1 \\ & & & -1 & 2 & -1 \\ & & & & -1 & 1 & \\ & & & & & & \ddots \end{pmatrix} \quad (14)$$

in which case the interblock Hamiltonian H_{BB} which describes the interaction between blocks becomes:

$$H_{BB} = \begin{pmatrix} 0 & & & & & \\ & 0 & & & & \\ & & 1 & -1 & & \\ & & -1 & 1 & & \\ & & & & 0 & \\ & & & & & 1 & -1 \\ & & & & & -1 & 1 & \\ & & & & & & & \ddots \end{pmatrix} \quad (15)$$

H_{BB} in turn also coincides with the incidence matrix of a graph which contains the links missing in Fig.4 (RHS) which connects consecutive blocks. In a few words: our RG-prescription introduces free BC's at the ends of every block. This condition fixes uniquely the breaking of H_{Free} into the sum $H_B + H_{BB}$. This is the choice we make. It should be emphasized that the splitting of H_{Free} into two parts $H_B + H_{BB}$ is by no means unique, so that different choices may lead to very different results. Prior to any computation we notice that the previous RG-prescription should lead to an exact value of the ground state energy, for the ground state of each block is again a constant function. The question is therefore to what extent our method is capable of describing the excited states. We shall concentrate ourselves to the first excited state since computations can be carried out analytically.

First of all we diagonalize H_B within each block of 3 sites, keeping only the ground state $\psi_0^{(0)}$ and the first excited state $\psi_1^{(0)}$ ($3 \rightarrow 2$ truncation). The superscript denotes the initial step in the truncation method. In the standard RG method we would choose $\psi_0^{(0)}$ and $\psi_1^{(0)}$ as the orthonormal basis for the truncated Hilbert space and obtain the effective Hamiltonian H'_B and H'_{BB} . In our case it is convenient to express these effective Hamiltonians in a basis expanded by the following linear combination:

$$\psi_+^{(0)} = \frac{1}{\sqrt{2}}(\psi_0^{(0)} + \psi_1^{(0)}) \quad (16)$$

$$\psi_-^{(0)} = \frac{1}{\sqrt{2}}(\psi_0^{(0)} - \psi_1^{(0)}) \quad (17)$$

which are also an orthonormal basis of the truncated Hilbert space. In this basis the truncation of H_B reads as follows,

$$H_B \longrightarrow H'_B = \begin{pmatrix} A & 0 & 0 & & \\ 0 & A & 0 & & \\ 0 & 0 & A & & \\ & & & \ddots & \\ & & & & A \end{pmatrix} \quad (18)$$

$$A = \frac{\epsilon}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

with ϵ taking on the value $\epsilon^{(0)} = 1$ in the initial step of the RG-method, which is the energy of the state $\psi_1^{(0)}$.

The truncation of H_{BB} is more complicated, the result being:

$$H_{BB} \longrightarrow H'_{BB} = \begin{pmatrix} B & C & 0 & & \\ C^t & D+B & C & & \\ 0 & C^t & D+B & & \\ & & & \ddots & \\ & & & D+B & C \\ & & & C^t & D \end{pmatrix} \quad (19)$$

$$B = \begin{pmatrix} a^2 & ab \\ ab & b^2 \end{pmatrix} \quad C = \begin{pmatrix} -ab & -a^2 \\ -b^2 & -ab \end{pmatrix} \quad D = \begin{pmatrix} b^2 & ab \\ ab & a^2 \end{pmatrix}$$

with a b taking on the values $a^{(0)} = \frac{1}{\sqrt{6}} - \frac{1}{2}$ and $b^{(0)} = \frac{1}{\sqrt{6}} + \frac{1}{2}$ in the initial step of the RG-method.

The nice feature about the basis (16)-(17) is that all rows and columns of (18) and (19) add up to zero, just like the original Hamiltonians (14) and (15), implying that the constant vector is an eigenvector with zero eigenvalue of the renormalized Hamiltonian!

We shall call $H_{N/3}(\epsilon, a, b)$ the sum of the Hamiltonians (18) and (19) for generic values of ϵ , a and b . Next step in our RG-procedure is to form blocks of 4 states of the new Hamiltonian $H_{N/3}(\epsilon, a, b)$ and truncating to the two lowest $\psi_0^{(1)}$ and $\psi_1^{(1)}$ energy states within each 4-block ($4 \rightarrow 2$ truncation). The reason for this change in the number of sites per block (from 3 to 4) is motivated by the form of H'_{BB} in (19) and the fact that if we try to make a second step in the RG-method with 3-blocks the method is doomed to failure because the constant state of Fig.2 would no longer be the ground state. Fortunately enough, with 4-blocks if we define new states $\psi_+^{(1)}$ and $\psi_-^{(1)}$ in the same form as we did in Eq.(5), we obtain that the new effective Hamiltonian is obtained by a redefinition of the parameters, namely,

$$H_{N/3}(\epsilon, a, b) \longrightarrow H_{N/6}(\epsilon', a', b') \quad (20)$$

$$\epsilon' = \frac{\epsilon}{2} + a^2 + b^2 - \Delta \quad (21)$$

$$a' = \frac{1}{2\sqrt{2}} \left[a + b - \frac{a(a^2 - 3b^2 + \Delta) + \frac{b\epsilon}{2}}{\sqrt{\Delta(\Delta + a^2 - b^2)}} \right] \quad (22)$$

$$b' = \frac{1}{2\sqrt{2}} \left[a + b + \frac{a(a^2 - 3b^2 + \Delta) + \frac{b\epsilon}{2}}{\sqrt{\Delta(\Delta + a^2 - b^2)}} \right] \quad (23)$$

$$\Delta \equiv \sqrt{(a^2 - b^2)^2 + \left(\frac{\epsilon}{2} - 2ab\right)^2} \quad (24)$$

In this fashion, the constant state of Fig.2 is again the ground state of the model and moreover, upon iteration of Eqs.(20)-(24) there are no level crossing among the excited states. Otherwise stated

this means that the level structure of the block Hamiltonian H_B is preserved under the action of our BRG-method based upon the reduction from 4 to 2 states.

The energy $E_1(N)$ of the first excited state of a chain with $N = 3 \times 2^m$ sites can be obtained iterating m times Eqs.(20)-(24):

$$E_1(N = 3 \times 2^m) \equiv \epsilon^{(m)} \quad (25)$$

The initial data are given by:

$$\epsilon^{(0)} = 1, \quad a^{(0)} = \frac{1}{\sqrt{6}} - \frac{1}{2}, \quad b^{(0)} = \frac{1}{\sqrt{6}} + \frac{1}{2} \quad (26)$$

In Table 2 we give our results for small and large number of iterations. For low values of N the deviation of $\epsilon^{(m)}$ with respect to the exact result is small. Recall that we are only keeping two states in our RG-procedure, and that the ground state energy is exactly zero by construction!. But what is more interesting about these results in Table 2 is that we are able to obtain the correct size dependence, i.e., $1/N^2$ of $\epsilon^{(m)}$. As a matter of fact, the energy of the first excited state behaves for large N as (8):

$$E_1^{(exact)}(N) \sim c_{exact}/N^2, \quad \text{with } c_{exact} = \pi^2 \quad (27)$$

while our BRG-method gives,

$$E_1^{(BRG)}(N) \sim c_{BRG}/N^2, \quad \text{with } c_{BRG} = 12.6751 \quad (28)$$

The achievement of the $1/N^2$ -law is a remarkable result which in turn allows us to match the correct order of magnitude of the energy. For instance, for 10 iterations our RG-method with 2 states kept gives the energy of the order of 10^{-6} , which is precisely the same order of magnitude as for the CBC method (see Table 1) but with 8 states kept in the case of Fixed BC's. Recall that the standard BRG performs as bad as a 10^{-2} order of magnitude.

3 Wave-Function Reconstruction

We may also wonder whether we are able to make a reasonable picture of the first excited state wave-function based upon our BRG-procedure when compared with the exact form depicted in Fig.2. As we are working with a real-space realization of the renormalization group method, this is something we have at hand. To do this we need to perform a “reconstruction” of the wave-function. This reconstruction amounts to plot the form of our aproximate wave-function in each and every of the 3-blocks out of the 2^{m+1} in which the original chain is decomposed into under the BRG-procedure. Recall that in the initial step we started out with blocks of 3 states keeping the two lowest states $\psi_0^{(0)}$ and $\psi_1^{(0)}$ ($3 \rightarrow 2$ truncation). In the next step we make blocks of 4 states keeping the two lowest states $\psi_0^{(1)}$ and $\psi_1^{(1)}$ ($4 \rightarrow 2$ truncation) and then we perform the iteration procedure over and over. As a result of this procedure we may express the two lowest wave functions of the $m+1$ -th step in terms of those of the previous m -th step by means of the following matricial form:

$$\begin{pmatrix} \psi_0^{(m+1)} \\ \psi_1^{(m+1)} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ \alpha_m & \beta_m \end{pmatrix} \begin{pmatrix} \psi_0^{(m)} \\ \psi_1^{(m)} \end{pmatrix}_L + \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -\alpha_m & \beta_m \end{pmatrix} \begin{pmatrix} \psi_0^{(m)} \\ \psi_1^{(m)} \end{pmatrix}_R \quad (29)$$

where the LHS of Eq. (29) represents the wave function of $3 \times 2^{m+1}$ sites while in the RHS we have a left-wave-function of 3×2^m sites and another right-wave-function of 3×2^m sites, so that everything squares. The parameters appearing in Eq. (29) turn out to be given by:

$$\alpha_m = \frac{(\frac{\epsilon_m}{2} - 2a_m b_m) + (a_m^2 - b_m^2 + \Delta_m)}{2\sqrt{\Delta_m(\Delta_m + a_m^2 - b_m^2)}} \quad (30)$$

$$\beta_m = \frac{(\frac{\epsilon_m}{2} - 2a_m b_m) - (a_m^2 - b_m^2 + \Delta_m)}{2\sqrt{\Delta_m(\Delta_m + a_m^2 - b_m^2)}} \quad (31)$$

with Δ_m as in Eq.(24). Their initial values are $\alpha_0 = 1/\sqrt{10}$ and $\beta_0 = 3/\sqrt{10}$. We may recast Eq. (29) in more compact form by writing:

$$\Psi^{(m+1)} = L_m \Psi_L^{(m)} + R_m \Psi_R^{(m)} \quad (32)$$

where

$$L_m = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ \alpha_m & \beta_m \end{pmatrix} \quad (33)$$

$$R_m = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -\alpha_m & \beta_m \end{pmatrix} \quad (34)$$

We may call Eq.(32) the *reconstruction equation*. This is the master equation that when iterated “downwards” (reconstruction) allows us to obtain the picture of our approximate BRG-wave-function corresponding to every and each block of 3 sites of the 2^{m+1} blocks in which the chain is decomposed into. At the end of the iteration procedure we end up with expressions for the values of the 3-sites wave-functions in terms of the initial two lowest states $\psi_0^{(0)}$ and $\psi_1^{(0)}$. The first one is a constant function while the second is a straight line of negative slope. Thus, these two states turn out to be the building blocks of our BRG-procedure. As a matter of illustration, we present in Fig.5 the plot for the reconstruction of the first excited wave-function for a chain of $N = 48$ ($m=3$) sites long. We can observe that our BRG-wave-function has the shape of a broken line around the exact form of the wave function. We have checked that this broken shape remains when increasing the length of the chain so that its nature must be due to the fact that we are only keeping 2 lowest states in our procedure. It is expectable that upon increasing this number the shape of the approximate wave function must be smoothed out. Nevertheless, the important point is that our BRG-method preserves the number of nodes of the wave-function which one of the defining features of the exact result.

When using the reconstruction equation to obtain the wave function we may use a binary code based upon the labels L (left) and R (right) to keep track of the different 3-sites blocks which make up the chain. Thus, in one dimension the RG-blocks are in a one-to-one correspondence with a binary numerical system. In general, for other dimensions we may state schematically the following correspondence:

$$\text{BRG-prescription} \longleftrightarrow \text{“Number System”}$$

4 Conclusions and Outlook

In this letter we have shown that it is possible to perform a successful real-space block RG treatment of the 1D tight-binding model. In doing so we have elucidated the role played by the boundary conditions in the real-space RG method for we have shown that the failure of the BRG-method pointed out in [9] can be traced back to a particular choice of boundary conditions: if we choose free BC’s instead of fixed BC’s the ground state turns out to be homogeneous, a property which is necessary in order to have a suitable implementation of a blocking procedure. Otherwise, for fixed BC’s the blocks of the RG-method cannot have fixed BC’s as well for in that case one introduces anomalous behaviour in the

ground state wave-function which ruins the method. Thus, the failure of the BRG-method accepted so far is not intrinsic to the method but depends on the choice of the boundary conditions.

We have devised a BRG-method which always yields the correct ground state of the Hamiltonian with free BC's for open chains of any length. As for the excited states, our method is able to reproduce the correct $1/N^2$ -law exhibited by the exact first excited state (27)-(28) in the large N -limit with the correct order of magnitud. Since we only keep 2 states - to make computations analytical - at every step of our truncation procedure, this means that we have been able to capture the essential physics of the model in the most economic way. For this reason we believe that our method should also be able to account for the correct properties of the rest of excited states of the model.

We have given another physical interpretation for the tight-binding Hamiltonian with free BC's as the restriction of the Ferromagnetic Heisenberg model to the subspace of one-magnon solutions (13). In the light of this mapping we may interpret the $1/N^2$ -law of our BRG-method as in good agreement with the correct *quadratic* dispersion relation exhibited by the Ferromagnetic magnons $E(k) \sim k^2 \sim 1/N^2$ for low wave number k . Thus, from this point of view, our RG-procedure is aplicable to the Ferromagnetic regime of the Heisenberg model and we may wonder whether we may extend it to the Antiferromagnetic regime and obtain the *linear* dispersion relation $E(k) \sim k \sim 1/N$.

Our BRG-method also gives a good qualitative picture of the ground state and first excited states wave-functions in coordinate space as shown in Fig.5. In fact, the number of nodes is preserved by the BRG-procedure. This good performance is due to a reasonable choice of the 2 lowest energy states kept in the initial stage of the method. The reconstruction of the wave-function is based upon the master equation (32) which can be readily generalized to account for a bigger number of states kept.

Finally, we may address yet another generalization of the present work by considering a Hamiltonian $H_{Free}(q)$ as in Eq.(6) but depending upon a complex parameter q . This parameter enters in the matrix expression of the Hamiltonian by replacing the diagonal $(1, 2, 2, \dots, 2, 1)$ of Eq.(6) by $(q, q + q^{-1}, q + q^{-1}, \dots, q + q^{-1}, q^{-1})$ while the off-diagonal terms remain as -1. This corresponds to a one-parameter deformation of the boundary conditions in which the free case is recovered as the $q \rightarrow 1$ limit. Furthermore, it is more interesting to consider this extension from the point of view of the Heisenberg-model connection for in this case the Hamiltonian comes with a boundary term depending on the q parameter:

$$H_\lambda = -\frac{1}{2} \sum_{j=1}^{N-1} (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \frac{q + q^{-1}}{2} (\sigma_j^z \sigma_{j+1}^z - 1)) - \frac{q - q^{-1}}{4} (\sigma_1^z - \sigma_N^z) \quad (35)$$

The important point about this Hamiltonian is that it is invariant under what is called the quantum group $SU_q(2)$. We have shown in [12] that this symmetry plays an important role in constructing an improved version of the standard BRG-procedure. We call this new method a q -BRG-method and for instance it is able to obtain correctly the line of critical XXZ models, unlike the standard BRG-method. This quantum symmetry must show up in the free case when performing the previous q -deformation.

e-mail addresses: mardel@fis.ucm.es (M.A. M.-D.); sierra@cc.csic.es (G.S.)

Table captions

Table 1 : Exact, Standard RG and CBC Values of Low Lying States for the 1D Tight-Binding Model for a chain of $N = 2048$ sites.

Table 2 : Exact and new standard RG values of the first excited state for the 1D Tight-Binding Model.

Figure captions

Figure 1: Ground state ψ_0 and first excited state ψ_1 for the Hamiltonian H_{Fixed} with fixed BC's.

Figure 2 : Ground state ψ_0 and first excited state ψ_1 for the Hamiltonian H_{Free} with fixed BC's.

Figure 3 : The one-dimensional open chain as a graph.

Figures 4 : Lattice decomposition into blocks of n_s sites ($n_s = 3$).

Figure 5 : The reconstruction of the first excited wave-function for a chain of $N = 48$ ($m=3$) sites long.

References

- [1] K.R. Wilson, *Rev. Mod. Phys.* **47**, 773 (1975).
- [2] Hirsch, J. 1983, *Phys. Rev. B* **28**, 4059, 1985, *Phys. Rev. B* **31**, 4403; Hirsch, J. and Lin, H.Q., 1988, *Phys. Rev. B* **37**, 5070.
- [3] S.D. Drell, M. Weinstein, S. Yankielowicz, *Phys. Rev. D* **16**, 1769 (1977).
- [4] R. Jullien, P. Pfeuty, J.N. Fields, S. Doniach, *Phys. Rev. B* **18**, 3568 (1978).
- [5] Hirsch, J. 1980, *Phys. Rev. B* **22**, 5259.
- [6] J.M. Rabin, *Phys. Rev. B* **21**, 2027 (1980).
- [7] Pfeuty, P., Jullien, R. and Penson, K.A., in “Real-Space Renormalization”, editors Burkhardt, T.W. and van Leeuwen, J.M.J., series topics in Current Physics **30**, Springer-Verlag 1982.
- [8] González, J., Martín-Delgado, M.A., Sierra, G., Vozmediano, A.H., *Quantum Electron Liquids and Hight- T_c Superconductivity*, Lecture Notes in Physics, Springer-Verlag 1995, to be published.
- [9] S.R. White, R.M. Noack, *Phys. Rev. Lett.* **68**, 3487 (1992).
- [10] Wilson, K.G., 1986, *unpublished informal talk*.
- [11] S.R. White, *Phys. Rev. Lett.* **69**, 2863 (1992); *Phys. Rev. B* **48**, 10345 (1993).
- [12] Martín-Delgado, M.A., Sierra, G., 1995, *UCM-CSIC preprint, to be published*.

Energies	Exact	Standard RG	CBC
E_0	2.351×10^{-6}	1.9207×10^{-2}	2.351×10^{-6}
E_1	9.403×10^{-6}	1.9209×10^{-2}	9.403×10^{-6}
E_2	2.116×10^{-5}	1.9214×10^{-2}	2.116×10^{-5}

Table 1: Exact, Standard RG and CBC Values of Low Lying States for the 1D Tight-Binding Model for a chain of $N = 2048$ sites.

m	N	$E_1^{(exact)}(N)$	$\epsilon^{(m)}$
0	3	1	1
1	6	0.2679	0.2792
2	12	0.0681	0.0765
10	3072	1.0458×10^{-6}	1.3421×10^{-6}
20	$\sim 3 \times 10^6$	9.9737×10^{-13}	1.2809×10^{-12}
32	$\sim 10^{10}$	5.9448×10^{-20}	7.6347×10^{-20}
	$\gg 1$	π^2/N^2	$12.6751/N^2$

Table 2: Exact and new standard RG values of the first excited state for the 1D Tight-Binding Model.